FORM PTO-1390 U.S. DE	PARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	Transmin nagranian and			
(REV 10-94)		7885.55USWO			
	TO THE UNITED STATES	7883.33USWU			
DESIGNATED/ELECTE	` '				
CONCERNING A FILIN	G UNDER 35 U.S.C. 371				
		US APPLICATION NO (If known, see 37 C F R 1 5) 09/125751			
INTERNATIONAL APPLICATION NO.	INTERNATIONAL FILING DATE	PRIORITY DATE CLAIMED			
PCT/NO97/00074	March 12, 1997	March 13, 1996			
TITLE OF INVENTION METHOD OF KILLING TARGET CELLS I	OTTIADARGEED OFFI DODIN ATIONS WA				
APPLICANT(S) FOR DO/EO/US	IN HARVESTED CELL POPULATIONS WI	TH ONE OR MORE IMMUNOTOXINS			
Oystein FODSTAD; Gunnar KVALHEIM; S.	_				
Applicant herewith submits to the United States De	esignated/Elected Office (DO/EO/US) the following	g items and other information:			
[X] This is a <b>FIRST</b> submission of items con-	cerning a filing under 35 U.S.C. 371.				
2. [ ] This is a SECOND or SUBSEQUE	NT submission of items concerning a filing under 3	35 U.S.C. 371.			
3. [X] This express request to begin national examination until the expiration of the approximation of the approximation until the expiration until the expi	amination procedures (35 U.S.C. 371(f)) at any time oplicable time limit set in 35 U.S.C. 371(b) and PCT	e rather than delay			
4. [X] A proper Demand for International Prelin	ninary Examination was made by the 19th month fr	Articles 22 and 39(1).  com the earliest claimed priority date.			
5 [X] A copy of the International Application at a. [X] is transmitted herewith (required	is filed (35 U.S.C. 371(c)(2))  If not transmitted by the International Bureau	n			
b. [X] has been transmitted by the Inter	rnational Bureau.				
a. [X] is transmitted herewith (required by the Interest control of the International Application of the Internation of the International Application of the International Application of the Internat	cation was filed in the United States Receiving Offi plication into English (35 U.S.C. 371(c)(2)).	ice (RO/US)			
a. [ ] are transmitted herewith (re	a. [ ] are transmitted herewith (required only if not transmitted by the International Bureau).				
b. [ ] have been transmitted by the International Bureau.					
c. [ ] have not been made; nower		NOT expired.			
anh upp					
8 [ ] A translation of the amendments to the	he claims under PCT Article 19 (35 U.S.C. 371(c)(3	3)).			
fine control of the c	[X] An unsigned oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).				
10 A translation of the annexes to the In	nternational Preliminary Examination Report under	PCT Article 36			
(35 U.S.C. 371(c)(5)).					
Items 11. to 16. below concern document(s) or in					
11. [ ] An Information Disclosure Statement					
12. [ ] An assignment document for recording	ng. A separate cover sheet in compliance with 37 C	FR 3.28 and 3.31 is included.			
13. [X] A FIRST preliminary amendment.					
[ ] A SECOND of SUBSEQUENT preli	iminary amendment.				
14. [ ] A substitute specification.					
15. [ ] A change of power of attorney and/or	r address letter.				
16. [X] Other items or information: International	Preliminary Examination Report				
		!			

300 Rec 0 PUI/PIU 25 AUG 1930

09/125751

U.S. APPLICATION NO (If know	n, see 37 C F R 15)	PCT/NO97/00074	4O	7885.55USWO	
17. [X] The following	fees are submitted:	<u> </u>		CALCULATIONS PT	O USE ONLY
BASIC NATIONAL FI	EE (37 CFR 1.492(a) (1)-(	5)): or JPO	\$930.00		O USE ONE
	inary examination fee paid	to USPTO	\$720.00		
		aid to USPTO (37 CFR 1.487 CFR 1.445(a)(2))			
	al preliminary examination in fee (37 CFR 1.445(a)(3)) p	fee (37 CFR 1.482) nor paid to USPTO	\$1,070.00		
	inary examination fee paid fied provisions of PCT Arti	to USPTO (37 CFR 1.482) cle 33(2)-(4)	\$98.00		
	ENTER APPROP	PRIATE BASIC FEE	AMOUNT =	\$1,070.00	
	r furnishing the oath or dec claimed priority date (37 C	laration later than [] 20 [] 3 FR 1.492(e)).	30	\$	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	12 -20=	0	X \$22.00	\$0	
Independent claims	1 -3 =	0	X \$82.00	\$0	
MULTIPLE DEPENDE	NT CLAIM(S) (if applicab	le)	+ \$270.00	\$	
	TOTAL	OF ABOVE CALCU	LATIONS =	\$1,070.00	
	ng by small entity, if applic filed (Note 37 CFR 1.9, 1.2)	able. Verified Small Entity 7, 1.28).		s	
<u>्र</u> ्थ		S	UBTOTAL =	\$1,070.00	
Processing fee of \$130.00 for furnishing the English translation later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(f).					
		TOTAL NATIO	NAL FEE =	\$1,070.00	
TOTAL NATIONAL FEE = \$1,070.00  Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property + \$					
		TOTAL FEES E			
				Amount to be:	\$
charged \$					
a. [X] A check in the amount of \$ 1,070.00 to cover the above fees is enclosed.  b. [] Please charge my Deposit Account No					
A duplicate copy of this sheet is enclosed.					
c. [X] The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 13-2725. A duplicate copy of this sheet is enclosed.					
NOTE: Where an app 1.137(a) or (b)) must be	propriate time limit under e filed and granted to rest	37 CFR 1.494 or 1.495 has ore the application to pend	s not been met, a ling status.	petition to revive (37 CFF	
send all correspondence Michael B. Lasky				Muly	WM
MERCHANT & GOT 3100 Norwest Center				SIGNATURE. (	
90 South Seventh Str	reet			Michael B.	Lasky
Minneapolis, MN 55	403			NAME	1
				29,555 REGISTRATION NU	MBER

# IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant:

FODSTAD et al.

Docket No.:

7885.55USWO

Serial No.:

Unknown

Filed:

Int'l Appln No.:

PCT/NO97/00074

Int'l Filing Date:

March 12, 1997

Title:

METHOD OF KILLING TARGET CELLS IN HARVESTED CELL

POPULATIONS WITH ONE OR MORE IMMUNOTOXINS

**CERTIFICATE UNDER 37 CFR 1.10:** 

"Express Mail" mailing label number: EL039318435US

Date of Deposit: August 25, 1998

I hereby certify that this correspondence is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Assistant Commissioner for Patents, Washington, D.C. 20231.

By:

\_ M.L. Gren

Name: Mark Green

#### PRELIMINARY AMENDMENT

**Box PCT** 

**Assistant Commissioner for Patents** 

Washington, D. C. 20231

Dear Sir:

In connection with the above-identified application filed herewith:

#### IN THE ABSTRACT

Insert the attached Abstract page into the application as the last page thereof.

## IN THE SPECIFICATION

A courtesy copy of the originally-filed PCT specification is enclosed herewith, but the World Intellectual Property Office (WIPO) copy should be relied upon if it is already in the U.S. Patent Office.

#### IN THE CLAIMS

Applicant requests that the claims be entered as published in the International Preliminary Examination Report.

In the International Preliminary Examination Report claims, please amend the claims as follows:

- 3. [Twice amended] Method according to claim[s] 1, [2,] characterized in that the used antibodies are MOC31 and BM7, or fragments thereof.
- 4. [Twice amended] Method according to claim[s] 1, [2,] characterized in that the used antibodies are MOC31 and BM2 or 12H12, or fragments thereof.
- 5. [Twice amended] Method according to claim[s] 1, [2,] characterized in that the used antibodies are MOC31 and 595A6, or fragments thereof.
- 9. [Twice amended] A preparation of two [I]immunotoxins to kill cells according to the method in claim 1, **characterized in** that it contains two immunotoxins directed to antigens present on malignant cells.
- 10. [Twice amended] A preparation [Immunotoxins] according to claim 9, characterized in that the antibodies are selected among MOC31, and BM7, 595, BM2, 12H12 or combinations of these or fragments thereof and the toxin is native or recombinant Pseudomonas exotoxin A, or fragments thereof.
- 11. [Twice amended] Use of [the] a preparation of two immunotoxins according to claim 9, to produce a therapeutic agent against cancer.

Please renumber Claim 13 as Claim 12.

#### REMARKS

The above preliminary amendment is made to remove multiple dependencies from claims 3 to 5.

A new abstract page is supplied to conform to that appearing on the publication page of the WIPO application, but the new Abstract is typed on a separate page as required by U.S. practice.

Applicant respectfully requests that the preliminary amendment described herein be entered into the record prior to calculation of the filing fee and prior to examination and consideration of the above-identified application.

If a telephone conference would be helpful in resolving any issues concerning this communication, please contact Applicant's primary attorney-of record, Michael B. Lasky (Reg. No. 29,555), at (612) 336-4634.

Respectfully submitted,

MERCHANT, GOULD, SMITH, EDELL,

WELTER, & SCHMIDT, P.A.

3100 Norwest Center 90 South Seventh Street

Minneapolis, Minnesota 55 (612) 332-5309

Dated: August 25, 1998

Michael B. Lasky Reg. Nol 29,555

MBL/sef

PCT/NO97/00074 Intl Filing Date March 12, 1997 Docket No. 7885.55USWO

## **Abstract**

A method for killing unwanted target cells in a cell population comprising nucleated cells harvested from peripheral blood, or CD-34+ or similar early progenitor cells selected from the above nucleated cells or from bone marrow aspirates, in which method the cell population is in vitro or in vivo exposed to two or more immunotoxins selectively killing the malignant cells, is described. Furthermore the invention relates to the mixture of immunotixins, the use of the mixture and a kit for performing the method.

10

35

# PTOPCT Resid 2 5 AUG 1998

1410 DE2 09/125751

# Method for killing unwanted Target Cells

The present invention relates to selective purging of a cell population for target cells by exposing the cell population to a combination of two immunotoxins.

So called autologous stem cell transplants comprise isolated cells from blood or bone marrow from cancer patients which after pretreating the patients contain adequate amounts of immature progenitor blood and immune cells to restore the function of a non-functioning bone marrow shortly, or over a longer period of time after reinstalling the cells in the blood of the patients from which the cells are harvested.

Purging of autologous haematopoietic transplants using antibodies is known in the art when the transplants represent unselected bone marrow samples. Such a purging is published i.a. by Myklebust, A.T., Godal, A., Juell, S. and Fodstad, Ø. "Comparison of two antibody-based methods for elimination of breast cancer cells from human bone marrow". Cancer Res. (USA) 1994, 54/1 (209-214) and Myklebust, A.T., Godal, A., Pharo, A., Juell, S. and Fodstad, Ø. "Eradication of small cell lung cancer cells from human bone marrow with immunotoxins", Cancer Res. (USA) 1993, 53(16), 3784-88. Both publications use immunotoxins in which the antibody is conjugated to a toxin. The principle is to kill malignant cells from the harvested bone marrow cells before reinjection of the cell suspension into the patient.

In the recent years it is developed methods in which the principle actually is the opposite. Using the so called stem cell transplantation one tries to positively select from blood or bone marrow a subgroup of normal cells which are able to restore normal bone marrow function after the cells are reinstalled in the patient. These "stem cells" consist of a mixture of the most immature precursors for blood and immune cells and also more differentiated cells. The harvest of such cells can be performed either by so called apharesis of peripheral blood, a procedure taking one or more days, or by immuno-absorption/selection of CD34+ cells (immature progenitor cells) from blood or bone marrow using different techniques known in the art.

Stray, K.M. et al., "Purging tumor cells from bone marrow or peripheral blood using avidin, biotin, immuno adsorption" In: P.G. Adrian, G. Samuel and A.W-W. Diana (Eds), Advances in bone marrow purging and processing, pp. 97-103,

AMENDED SHEET

10

15

20

25

30

35

Orlando: Willelis Inc., 1994 describes purging of bone marrow cells or an apharesis product from peripheral blood. This procedure is performed for purging in patients with lymphoma and in patients with breast cancer. The method includes an enrichment step for CD34+ cells before purification of B-cells or breast cancer cells with a so-called avidin column. In this case the purging is performed indirectly in that the cell suspension initally is incubated with primary antibodies which bind to the breast cancer cells, the cell suspension is washed and once more incubated with an antibody which bind to the primary antibodies. This rat antibody is biotinylated, i.e. connected to a molecule which binds strongly to avidin. When this cell suspension finally is loaded on a column with avidin conjugated beads the tumor cells are trapped by the binding between cells with primary antibodybiotinylated secondary antibody-avidin. The results of purging using such a system was at the most 3.2 log depletion of malignant cells. The principle is very timeconsuming and cumbersome since the cell suspension must be handled in several steps including incubation with antibody and two washes before it is loaded on to the column. Thus, it is difficult to avoid damage to the stem cells or that there is unspecific trapping of the stem cells in the column, resulting in unfortunate loss of cells crucial for the recovery of normal bone marrow function.

Tyer, C.L. et al., "Breast cancer cells are effectively purged from peripheral blood progenitor cells using an immunomagnetic technique". Abstract to the first meeting of International Society for Hematotherapy and Graft Engineering, Orlando FL, 1993 describes an immunomagnetic method similar to the one used in the publication of Myklebust et al. above. This method is, however, used on peripheral blood cells. This principle is also entirely different from the use of immunotoxins and the effectivity of the purging varied from 3.3 to 4.8 log depletion of malignant cells in model experiments. The abstract does not mention any additional use of an indirect system with incubation of primary antibodies followed by washing and new incubation with beads connected to antibodies which bind to the primary antibody, but this is a reasonable assumption. Also this procedure comprises additional and possibly traumatic treatment of the normal cells and the procedure is time requiring. The effectivity is limited and the abstract does not mention anything about purging of CD34+ cell populations which will be a major problem with this method, as selection of CD34+ cells per se is time requiring and laborious. Thus, in most cases an immunomagnetic principle is used for CD34+ cell selection, which in this example is followed by one or two immunomagnetic steps for purging purposes. Therefore, there is a considerable risk for cell destruction and cell loss, with a method requiring a long lasting procedure and

involving high costs.

5

10

15

20

25

30

35

Lemoli, R.M. et al. (1994) Bone Marrow Transplant 13: 465-471 describe purification of human CD34<sup>+</sup> hematopoietic cells using the avidin-biotin immunoabsorption technique. They increased the purging of neoplastic cells by using several immunotoxins containing the ribosome-inactivating protein saporin and directed toward the lymphoid associated antigens CD30 and CD2. Tecce, R. et al (1991) Int. J. Cancer, 49: 310-316 describe purging autologous bone marrow prior to transplantation in patients suffering from monocytic leukemia with 2 monocytic-cell-lineage-specific immunotoxins constructed woth saporin and 2 MoAbs of high specificity for circulating monocytes and M5b acute nonlymphoid leukemia (ANLL). In WO91/09058 it is described immunotoxins comprising the myelomonocytic specific MoAb 195 useful for purging ANLL from bone marrow. Tonevitsky, A.G. et al (1986) Int. J. Cancer, 37: 263-273, describe purging of murine erythroleucemic stem cells from bone marrow employing an immunotoxin comprising a conjugate of ricin-A-chain and MoAb MAE15 which binds to the surface of normal and neoplastic murine erythroid cells: a model for studies of bone marrow transplantation therapy.

In isolating stem cells for transplantation one of the main objectives was that the cells which is intended to be reinstalled in the patients are selectively isolated in such a way that the transplants should not contain any malignant cells. In prior art it was recently shown that such preparations of stem cells surprisingly comprises malignant cells in a significant number of the examined cases. Up to now, very limited efforts have been made to remove or kill selected malignant cells in such transplants. This is partly because the skilled person of the art has not seen the need, and also because it was expected that the actual known methods were not specific and thus also would kill or remove the vulnerable stem cells. Furthermore, a supply of bone marrow or mobilized peripheral blood from a patient is not simple and unlimited and such a method for purging of stem cell transplants has to be performed within a short period of time and must be uncomplicated in order to avoid loss of, or damage to the normal cells can. Thus, referring to the above, the reason for using stem cell transplants is partly that the transplant should be completely free of cancer cells, partly that reconstitution of bone marrow function is faster than after transplantation with unselected bone marrow. It follows that it was absolutely necessary to invent a method which leave the fragile normal stem cells intact and which is practical to perform in combination with stem cell isolation procedures.

15

20

The object of the present invention is therefore to provide a method for purging of stem cell transplants which do not comprise the above disadvantages.

Such objects are obtained by the present invention characterized by the enclosed claims.

The present invention relates to purging of harvested stem cell populations in cases of solid tumors in which method the cell population is exposed to a composition of two or more antibodies connected to bacterial toxins. The used antibodies are directed to target cell-associated antigens.

In the following the invention is described in more detail by using an example of purging of stem cell transplants harvested from peripheral blood to remove breast cancer cells.

Known techniques to harvest cells comprise immuno-adsorption/selection of peripheral blood stem cells (PBSC) or CD-34<sup>+</sup> cells from blood or bone marrow. However, there are no known harmless and sufficiently effective techniques for purging of these cell populations for tumor cells. It seems evident that even among these immature cells there are malignant cells which according to prior knowledge should not posess CD34 receptors. Importantly, the invention described below surprisingly purged also the cancer cells without toxicity to normal progenitors.

Before harvesting a stem cell transplant from peripheral blood it is necessary to 25 mobilize the stem cells from the bone marrow by using chemotherapy or treatment with growth factors by methods known in the art. The harvesting of stem cells can be performed according to one or several methods, dependent on what kind of cells is desired. In one method peripheral blood stem cells are collected. This can be performed by scheduling patients to undergo leukapheresis on days 10 and 11 of 30 G-CSF administration (10 µg/kg/day) after receiving high doses of chemotherapy and total body irradiation. Blood flow rate may be fixed at for example 70 ml/min by using a CS-300 Plus blood cell separator (Baxter Healthcare Corporation, Fenwal Division, Deerfield, IL, USA). The average volume of blood treated during such a procedure may be about 10 liters for 21/2 hour to a dual-lumen central 35 venous catheter. Fifty ml of PBSC may be collected and washed with phosphate buffer saline (PBS), 1% human serum albumin (HSA) in a Cobe Processor 2991 to remove platelets. For use in the present invention the concentration of cells (2-4 x

10

15

30

 $10^{10}$ /apharesis) can be regulated to 1 x  $10^8$ /ml for negative selection (purging) with immunotoxins.

If CD-34<sup>+</sup> cells are desired this may be obtained by a positive selection with ISOLEX 50<sup>®</sup> or ISOLEX 300<sup>®</sup> (Baxter). In this method the product from apharesis or from bone marrow, which may be about 4 x 10<sup>10</sup> to 6 x 10<sup>10</sup> cells, can be mixed together and incubated with e.g. anti-CD34<sup>+</sup>-monoclonal antibody 9C5 at 0.5 µg/1 x 10<sup>6</sup> cells at 4°C for 30 minutes on a gentle rotator. The treated cells are washed with PBS with 1% HSA on Cobe Processor to remove unbound antibody. DYNAL<sup>®</sup> beads M-450 are added to the CD34<sup>+</sup> fraction at 0,5 beads per 1 nucleated cell at 4°C for 30 minutes. Magnetic separation of rosettes from non-targeted cells can be performed by washing away unbound cells two or three times with PBS, 1% HSA. The CD34<sup>+</sup> cells can then be released from the Dynal<sup>®</sup> beads by adding, for example ChymoCell-R (Chymopapain) at a final concentration of 200 pKat/ml in 15 minutes at room temperature. Thus CD34<sup>+</sup> cells can be harvested by washing with PBS in 5% sodium citrate. Also other procedures for selecting stem cells/early progenitor cells are known.

The binding profile of several antibodies in breast cancer cell lines and tumor
materials has been examined by others and partly confirmed by us. The antibodies
which bind to a large percent of breast cancer cells and not to important immature
normal cells in blood and bone marrow, were conjugated to Pseudomonas exotoxin
A (PE) and the ability to kill breast cancer cells in culture was examined, mainly in
colony producing assays. Based on the binding profile of the antibodies the present
inventors produced in all five different immunotoxins:

- 1. MOC31-PE: This conjugate binds to a very high percent of all breast cancer cells and was very effective in the model experiments and in the actual concentrations with marginal toxicty to normal hematopoietic progenitor cells.
- 2. NrLu10-PE: This binds to the same antigen as above, but to another epitope. It is slightly less active than MOC31-PE.
- 3. BM7-PE: This binds to the protein part of a mucin antigen which mainly is found on breast cancer cells. The antigen is present on a great percentage of breast cancer cells, but not all. The immunotoxin showed high specific activity to the cancer cells, but was not as effective as the two previous immunotoxins.

25

30

35

- 4. BM2-PE: This binds to a sugar containing epitope on the same antigen as BM7-PE. The immunotoxin showed approximately the same effectivity as BM7-PE concomitant with a very low toxicity for normal cells.
- 5 MLuC1-PE: This binds to a totally different antigen, the Lewisy-antigen. The immunotoxin was slightly less active than the previous one and also showed a moderate toxicity to normal cells.
- The immunotoxins according to 1, 2, 5 were tried individually and in combination in model experiments of purging regular bone marrow samples for cancer cells (Myklebust et al., Cancer Research, 1994). It is, as already mentioned, a great advantage to use stem cell transplants because of a shorter interval to regain normal bone marrow function (i.e. safer procedure). However, in spite of what was expected, such transplants are contaminated with tumor cells and it was necessary to apply immunotoxins able to kill <u>all</u> cancer cells in the transplant without significantly affecting the normal cells.

Search for immunotoxins more specific for breast cancer than MLuC1-PE and better suited for purging in breast cancer was initiated.

During this search it was surprisingly discovered that the combination of MOC31-PE and BM7-PE were more effective than the sum of each of the previous immunotoxins used alone. This is demonstrated in Table 1 below. Further studies of binding between the antibody and the cell lines have shown that the combination result in a stronger binding than the single antibodies.

MOC31 binds to most of the breast cancer cells, also those which are less differentiated. BM7 recognizes a mucin antigen which is expressed on a considerable fraction of breast cancer cells, also on the cells which are more differentiated. NrLu10 and BM2 bind to the antigen recognized by MOC31 and BM7, respectively, and with this in mind it was not very probable that they would add anything to a combination of MOC31 and BM7 immunotoxins. MLuC1-PE was theoretically interesting in that it is binding to a different antigen than the immunotoxins mentioned above. MLuC1-PE was however, toxic for normal cells and the model experiments did not show any clear advantage by including this in the combination.

In the example below, purging of peripheral stem cell transplants (apharesis

10

15

20

products) is described. In addition, the inventors have performed several experiments using the combination MOC31-PE and BM7-PE in experiments where tumor cells are added to harvested peripheral stem cells or bone marrow before an immunomagnetic positive selection of CD-34+ cells. The results from one such experiment is shown in Table 2. With two different cell lines it is demonstrated that the positive selection of CD34<sup>+</sup> cells in itself (without any other form of purging) removes up to 3.8 log tumor cells from the original harvested cell population. In other experiments the "purging" effect of CD34 selection varies from 2-3 log which also is referred to in the literature. When the immunotoxin treatment was used on the positively selected CD34 population the total purging effect was more than 4.7 log (Table 2) for both cell lines. More than 4.7 log means in this case that all detectable tumor cells were removed. In other experiments we have in separate assays grown tumor cells and normal progenitor cells taken from CD34+ population after 1 hour treatment with immunotoxin. In these experiments we observed that the tumor cells are killed or dying shortly after treatment while the tumor cells in unpurged control populations grew and created colonies, and/or proliferated in cell-adherence type cultures. The normal stem cells are not influenced by the treatment such that in three different test systems the survival of the normal progenitor cells are only insignificantly reduced. Table 3 shows a similar experiment in which the CD34+ cells were incubated with the immunotoxins for 2 hours at 37°C, and it is demonstrated that the stem cells essentially survive the immunotoxin treatment.

Table 1 Effect of immunotoxins involving PE in killing PM1 breast cancer cells

			A CONTRACTOR OF THE CONTRACTOR	The second secon
PE-Immunotoxin with MAbs	Exp	Log cell ki	Log cell kill at Immunotoxin concentration of	ration of
		0.01 µg/ml	0,1 µg/ml	1,0 µg/ml
		mean ± s	mean ± s	mean ± s
BM2	3	$0.01 \pm 0.01$	$0.27 \pm 0.09$	$2.38 \pm 0.38$
BM7	3	$0.13 \pm 0.11$	$0.64 \pm 0.15$	$2.55 \pm 0.40$
MLuCI	3	$0.10 \pm 0.13$	$0.36 \pm 0.19$	$2.10 \pm 0.68$
M0C31	3	$0.81 \pm 0.10$	$2.83 \pm 0.29$	> 5
BM2+MOC31	4	$1.20 \pm 0.26$	> >	> 5
BM7+MOC31	4	$1.16 \pm 0.26$	> >	> 5
MLUC1+MOC31	4	$1.14 \pm 0.36$	> \$	> 5
BM2+BM7 MLuc1+MOC31	2	1.15 ± 0.2 - 1.55 ± 0.6	> 5	> 5

Table 2.

Purging effects of positive immunomagnetic selection of CD34<sup>+</sup> alone and combined with anti-breast carcinoma MOC31 and BM7 immunotoxin, in model experiments with PM1 and T-47D breast cancer cells admixed (1%) to peripheral blood stem cells.

Method		PM1			T-47D	
	No. of experiments performed	No. of cells depleted	Log cell killed	No. of experiments added	No. of cells depleted	Log cell killed
Positive selection of CD34 <sup>+</sup> cells	2	5 x 104	3.8 ± 0.04	2	5 x 104	$3.7 \pm 0.04$
Positive selection of CD34 <sup>+</sup> cells followed by IT treatment	2	5 x 104	> 4.7	7	5 x 10 <sup>4</sup>	> 4.7

a Calculated from observed number of colonies, taking into account the plating efficiency, of the number of cell killed by the treatment. b Mean of the results obtained in indpendent experiments, each performed in triplicate.

c MOC31-PE and BM7-PE immunotoxin were used at a concentration of 1 ug/ml of each.

Effect of IT on the survival of colonies in CD34<sup>+</sup> cells selected from mobilized peripheral blood. One x10<sup>5</sup> CD34<sup>+</sup> cells were incubated with the immunotoxins for 2 h at 37°C, seeded out in CFU-GM and CFU-GEMM (5 x 103/dish) cultures, and assayed as described in

"Material and Methods" in the Example.

		<del></del>	= =====================================	-
-GM	2º%	(100)	(77)	(53)
CFU-GM	No. of colonies mean ± SDb	85 + 6	97499	45 + 8
BFU-E	o%c	(100)	(68)	(98)
BFU	No. of colonies mean ± SD <sup>b</sup>	193 ± 31	172 ± 31	166 ± 14
ЕММ	2%	(100)	(95)	(06)
CFU-GEMM	No. of colonies mean ± SD <sup>b</sup>	456 ± 118	432 ± 56	$412 \pm 118$
Concentration of each IT (ug/ml) <sup>a</sup>			1.0	2.5
Treatment		None	MOC31-PE +BM7-PE	

It was very surprising that the use of two antibodies, directed to antigens expressed by epithelial cells according to the present invention, each combined with the Pseudomonas exotoxin A bacterial toxin, killed the malignant cells without performing any damage to the normal stem cells in the harvests from peripheral blood and bone marrow. It is known in the art that cells can be killed by bacterial exotoxins and that the killer effect is increased by connecting the toxin to antibodies directed to epitopes expressed by the target cells. However, it is also known that if immature cells are exposed to one or several immunotoxins it is a profound possibility that this treatment kills the normal stem cells in the cell population. Furthermore, these normal stem cells are sensitive to ex vivo treatment accompanying mechanical traumas and temperature changes. In the present invention the cell population, for example a stem cell transplant harvested from peripheral blood, is exposed to a composition of two antibodies each conjugated to PE. Since one of the immunotoxins was exceedingly active it is surprisingly demonstrated that by using a composition of two antibodies connected to a bacterial toxin the purging effect seemed greater than the sum of the effects when the immunotoxins are used separately. This synergistic effect is demonstrated in table 4 of the present disclosure using the antibodies BM7 and MOC31 connected to Pseudomonas exotoxin A in killing PM1 human breast cancer cells. The immunotoxins are both monoclonal antibodies directed against tumor associated antigens connected to the bacterial toxin Pseudomonas exotoxin A. One of the antibodies recognizes a epithelial antigen coded for by the GA 733-2 gene, which is expressed by most of the carcinoma cells and therefore can be used in all cases involving carcinomas (for example breast cancer, colorectal cancer, prostate cancer, ovarial cancer, lung cancer and pancreatic cancer). The other antibody is directed to mucin, a mucus protein which is slightly different from one carcinoma type to another. Commonly the antigen can be described as proteins encoded by the genes MUC-1, MUC-2 and MUC-3. Examples of the above mentioned monoclonal antibodies are MOC31 and BM7.

30

35

5

10

15

20

25

Conjugation of antibody and toxin can be performed in different known ways. Selection of two or more antibodies in the composition for linking to the bacterial exotoxins was performed in such a way that antibody binding was directed to epitopes expressed on a majority of the target cells and not on the normal cells. The problem in prior art is that both malignant cells and normal blood cells express common antigens on the cell surface. In the example enclosed in the present disclosure the two monoclonal antibodies MOC31 and BM7 are used. The former of these antibodies are directed to epithelial cells which if found in peripheral

blood, are malignant. The antigen (the whole protein) is encoded by the gene GA 733-2. This antigen has, however, several epitopes and it is important to target the epitopes most abundantly expressed.

The BM7 antibody is one of the antibodies directed to an epitope of the antigen expressed by the MUC1 gene. Several genes encode similar antigens e.g. (MUC2, MUC3).

The bacterial toxin Pseudomonas exotoxin A has a relatively moderate toxic effect on normal stem cells and malignant cells. However, when connected to antibodies directed to antigens expressed on the target cells the toxic effect on these is very pronounced. In Table 5 it is demonstrated that the mixture of immunotoxins according to the invention, even after an incubation time of as little as 60 minutes, kills T-47 D cells, MCF7 cells and PM1 cells at a much higher degree of efficacy than known in the prior art with other methods. The combination of these two immunotoxins is thus giving surprising results in relation to what should be anticipated because of the selective efficacy, simplicity and the only marginal toxicity to normal progenitor cells.

It may be claimed that it is known to use several immunotoxins consisting of Pseudomonas exotoxin A conjugated to three different antibodies, see Myklebust et al. 1993 and 1994. One of the antibodies (MOC31) were used in a similar way to the present invention in order to purge unselected bone marrow cells. However, the other antibodies used do not seem to be optimal, among other things because one of them is directed to the same antigen as MOC31 and because the other (MLuC1) crossreact with normal cells and thereby the immunotoxin linked to this antibody could easily be toxic for the most immature stem cells. In the present invention on purging stem cell preparations we have prepared another monoclonal antibody which add to the effect of MOC31 and observe that the combination of these two antibodies used as immunotoxins give profoundly surprising results.

Due to the high specific activity of the disclosed immunotoxins it seems possible to administer the mixture for in vivo treatment of patients suffering from different types of carcinoma. If the cancer disease is limited it will be possible to inject or infuse each of them or the mixture intravenously for example when spreading of the disease to the bone marrow is demonstrated. It is further possible to inject the immunotoxins alone or in combination in patients with further spread of the disease with abdominal fluid (ascites) or with pleural effusion. A third possibility is

to treat patients with spread of the cancer to the central nervous system. In this case the immunotoxins can be injected directly into the tumor tissue, into the spinal fluid or in the artery supplying blood to the brain.

It is not known to use these immunotoxins in vivo except that MOC31-PE has been used in a leptomeningeal tumor model for small cell lung cancer. BM7-PE is not described in the literature at all.

An important problem in using immunotoxins in vivo is that their half lives often are very short, i.e. the immunotoxins are broken down and removed from the blood before the concentration is adequately high in the tumor. In US PS 5 322 678, Morgan et al. has patented a modification of the antibody part of a immunotoxin in order to reduce the problem with short half life in vivo. The present inventor suggests similar modification of the toxin part, a procedure not previously performed or known.

# Example 1

Effective purging of breast cancer cells from peripheral blood stem cell harvests with immunotoxins.

Introduction

20

25

30

35

High dose chemoradiotherapy with autologous hematopoietic stem cell support is being used with increasing frequency to treat patients with a variety of malignancies (1, 2). In the cases where this approach is unsuccessful, the most common reason is relapse of the disease, rather than toxicity, infections, and lack of engraftment (3). Importantly, there is solid evidence that in patients receiving high dose treatment, reinfusion of autografts containing clonogenic tumor cells can contribute to relapse and influence outcome (4). Gene-marking studies of autografted cells have indicated that tumor cells remaining in reinfused bone marrow (BM) contributes to recurrence of the disease (5). This conclusion is further supported by results in patients with follicular lymphomas which indicate that efficient BM purging improves disease-free survival (6.)

Using sensitive immunocytochemical techniques tumor cell contamination can be observed in histologically normal bone marrow autografts in the range of 37-62% of breast cancer patients undergoing high dose treatment (7). Peripheral blood stem cell (PBSC) autografts collected by apharesis after pretreatment with hematopoietic growth factors and chemotherapy are used to an increasing extent in the belief that

these products will have a low probability of containing tumor cells. However, recently it has been found that although tumor cell involvement is less extensive in PBSC autografts compared to BM harvests, malignant cells are still frequently found in mobilized PBSC collections from breast cancer patients (4, 7). In addition, recent findings show that chemotherapy and/or growth factors may mobilize tumor cells into the peripheral blood in patients both with and without prior detectable cancer cells in bone marrow (7, 8), results that further increase the risk of tumor cell contamination in PBSC grafts.

To avoid reinfusion of malignant cells, in vitro purging of PBSC autografts of breast carcinoma cells may be needed. Here we report a practical and rapid purging method, demonstrating that a 60 min incubation procedure with ITs directly added to the apharesis product selectively kills more than 5 log of tumor cells.

#### 15 MATERIALS AND METHODS

Cell Line. The PM1 breast cancer cell line was established in our laboratory from ascitic fluid drawn from a patient with advanced disease. The MCF7 and T-47D cell lines were obtained from the American Type Culture Collection (Rockwille, MD) (ATCC HTB 22 and ATCC HTB 133 respectively). Cells were cultured at 37°C in a 5% CO2 atmosphere in RPMI 1640 medium (RPMI) supplemented with 10% heat inactivated fetal calf serum (FCS) and antibiotics (100 U/ml of penicillin, 100  $\mu$ g/ml of streptomycin). Medium and supplements were purchased from GIBCO (Paisley, UK).

25

30

35

20

5

Human Bone Marrow and Peripheral Blood Progenitor Cells. BM cells were obtained from healthy volunteer donors. The BM mononuclear cell (MNC) fraction was obtained by Lymphoprep<sup>®</sup> (Nycomed Pharma, Oslo, Norway) and washed twice in phosphate buffered saline (PBS) before being used in the experiments. PBSCs were prepared from non-Hodgkin lymphoma patients. To mobilize PBSCs the patients were pretreated with chemotherapy plus hematopoietic growth factors (G-CSF, Neupogen, Amgen/Hoffman-La Roche, Basel, Switzerland). Eleven to twelve days after chemotherapy, when the number of CD34<sup>+</sup> cells in the pheripheral blood is high, the stem cells were collected by the use of a CS-3000 Plus blood cell separator (Baxter Healthcare Corp., Fenwal Division, Deerfield, IL).

20

25

30

Toxin, Antibodies, and Construction of Immunotoxins. The anti-MUC1 (9) antibody BM7 (IgG1) was a gift from S. Kaul (Frauenklinik, University of Heidelberg, Germany), and the anti-EGP2 (10) antibody MOC-31 (IgG2a) was kindly provided by L. de Leij (University of Groningen, The Netherlands), and by MCA Development (Groningen). PE was obtained from Swiss Serum and Vaccine Institute (Bern, Switzerland). Each antibody was conjugated to PE via a thioether bond formed with sulfo-succinimidyl-4-(N-maleimidomethyl)cyclohexane-1-carboxylate (Pierce, Rockford, IL) as described previously (11).

Immunotoxin Treatment. The effect of IT treatment on clonogenic breast cancer cell survival was tested by incubating 2x10<sup>6</sup> exponentially growing tumor cells in RPMI with FCS with the indicated concentrations of ITs at 37°C with gentle agitation (100 rpm on an orbital incubator (Gallenkamp, Leicestershire, UK)) for varying periods of time, as indicated for each experiment. The cells were washed twice in PBS with 1% FCS before being seeded out in the clonogenic assay.

In some experiments, 10% tumor cells were admixed to BM mononuclear cells or PBSCs, incubated with ITs washed and the effect assessed for tumor cell or hematopietic progenitor clonogenic cell survival.

Colony Assays for Tumor and Hematopoietic Progenitor Cells. The clonogenic soft agar assay for tumor cells used has been described previously (12). Triplicate cultures were incubated for 14 days at 37°C in 5% CO<sub>2</sub>, 5% O<sub>2</sub>, and 90% N<sub>2</sub>, and colonies of more than 50 cells were counted in a Zeiss stereo microscope.

The clonogenic capacity of treated and untreated normal progenitor cells was assessed in CFU-GEMM assays (13) in which  $5x10^4$  PBSCs per ml were cultivated individually in standard methylcellulose cultures (HCC-4433 Methocult, Terry Fox Labs, Vancouver, BC) in IMDM medium (GIBCO) After 19 days of incubation, BFU-E and CFU-GM colonies were counted in an inverse phase contrast microscope. Each assay was performed with triplicate cultures in 1 ml 35 mm dishes at 37°C in a 5% CO<sub>2</sub>, 100% humidified atmosphere.

#### RESULTS

35

Growth of Human Breast Cancer Cells in Soft Agar. In several experiments, a linear relationship between the number of tumor cells seeded and the number of tumor colonies formed was observed. With the PM1 cell line the cloning efficiency

was in the range of 20 to 30% (not shown). In experiments with the T-47D and the MCF7 cell lines, the linear relationships previously reported (14) with PEs of 27% and 22%, respectively, were confirmed. These data were used to calculate efficiency of breast cancer cell depletion obtained with the treatment.

Efficacy of Individual and a Mixture of Immunotoxins in Killing Breast Cancer Cells. In model experiments three different concentrations of each IT were used. As demonstrated in Table 4, only marginal effects were obtained with the BM7 conjugate at the two lower concentrations, whereas 2.5 log cell kill was achieved at 1.0 μg/ml close to 3 log cell kill was seen, and at the highest concentration (1 μg/ml) the efficacy was at least 5 logs, the maximal effect possible to assess in this assay (14). With a mixture of both ITs, each at the indicated concentrations, all tumor cells were killed already at 0.1 μg/ml (Table 4). The results show that the mixture of the two ITs can kill the breast cancer cells very efficiently, and the data also suggest that additivity may be obtained by combining the two conjugates. Similar results were obtained when the efficacy of the ITs was tested against the two other breast cancer cell lines (not shown). Because of the expected heterogeneity in antigen expression on target cells, it seemed logical to use the IT combination in the further development of a method suitable for clinical use.

Efficacy of immunotoxins involving Pseudomonas exotoxin A in killing PM 1 human breast cancer cells. Table 4

PM1 cells were incubated with immunotoxins for 2 h at 370C, seeded out in soft agar, and colony formation was assessed as described in "Materials and Methods".

ITs with MAbs	Cell line	No. of	Log cell k	Log cell kill <sup>a</sup> at immunotoxin concentration of	tration of
		experi-			
		ments	A de la constantina del constantina de la constantina del constantina		
			0.01 µg/ml	0.1 µg/ml	1.0 μg/ml
			Mean <sup>b</sup> ± SD	Mean ± SD	Mean ± SD
BM7	PM 1	3	$0.13 \pm 0.11$	$0.64 \pm 0.15$	$2.55 \pm 0.40$
	MA 11		$0.96 \pm 0.18$	$2.02 \pm 0.07$	> 5
MOC-31	PM 1				
	MA 11	3	$0.81 \pm 0.10$	$2.83 \pm 0.29$	> 5
BM7+MOC-31c	PM I		$1.49 \pm 0.34$	$2.29 \pm 0.07$	> 5
	MA 11	4			
			$1.16 \pm 0.26$	> 5	> >
			$2.80 \pm 0.21$	> 5	> 5

a Calculated from observed number of colonies, taking into account the plating efficiency, and determined as logarithm of the number of cells killed by the treatment.

b Mean of the results obtained in independent soft agar experiments, each performed in triplicate.

c Each immunotoxin used at the concentration indicated.

10

Influence of Incubation Time. In the above experiments, a 120 min incubation with ITs was used. In a clinical setting it would, for practical reasons, be advantageous to use an even shorter incubation time. To study whether the time of exposure to the ITs might be reduced without affecting the specific tumor cell kill, the mixture of the two conjugates, used at a concentration of 1 µg/ml of each, was tested with different incubation times against the three breast cancer cell lines. In all cases, the 120 min exposure to the ITs resulted in eradication of all tumor cells. Importantly, the treatment was equally effective when the incubation time was reduced to 90 min and even to 60 min (Table 5), and the data demonstrate that at the IT concentrations used the shortest incubation time is sufficient for killing all clonogenic tumor cells present in the tumor cell cultures.

Influence of incubation time on the efficacy of a mixture of MOC-31 and BM7 immunotoxins<sup>a</sup> in killing breast cancer cells. Table 5

Tumor cells alone (2x106/ml) or admixed (ratio 1:10) to peripheral blood progenitor cells (total 1x107 cells/ml) were incubated with ITs at 370C for the indicated periods of time, seeded out in soft agar, and assayed as in Table 4.

Incubation time (min)		Mean log tumor cell kill with cell lineb	kill with cell lineb	
	T-47 D	MCF7	PM1	PM1 admixed to PBPCs
09	> 5	> 5	> 5	> 5
06	> 5	> >	> >	^
120	>5	> 5	> 5	> 5

a Each IT used at a concentration of 1 μg/ml.

b Result obtained in two independent experiments with T-47D and PM1 cells, and in one experiment with MCF7 cell.

To examine whether the toxicity to breast cancer cells might be altered in the presence of a high number of normal hematopoietic cells, experiments were performed in which the tumor cells were admixed at a ratio 1:10 to PBPCs harvested by apharesis. As is demonstrated in Table 5, the ITs killed more than 5 logs of PM1 tumor cells also in the presence of the normal cells, already after an incubation time of only 60 min. Since the results were the same for all three cell lines, the data indicate that the IT procedure may be effectively used in a clinical setting.

Influence of Incubation Conditions and Cell Concentration. To examine the efficacy of the IT procedure under conditions similar to those that would be used on clinical samples, PM1 tumor cells were admixed at a ratio of 1:10 to PBPCs in experiments in which unwashed cells taken directly from the apharesis bag before incubation with the ITs were resuspended in normal saline with ACD (stock solution bag, R 2220, Baxter Healthcare Corp., Fenwal Division). The results were compared to those obtained in the intial experiments with cells that were washed and resuspended in RPMI with 10% FCS. It was found (Table 6) that in both cases the treatment with the ITs for 60 min killed all PM1 cells, indicating that for clinical use the ITs can be injected directly into the apharesis bag, and that the low pH under such conditions does not affect the cytotoxicity of the ITs.

Effect of the incubation conditions on the efficacy of a mixture of MOC-31 and BM7 immunotoxins in killing PM1 breast cancer cells Table 6

admixed (ratio 1:10) to peripheral blood progenitor cells isolated by blood cell separator apharesis.

Cells from the apharesis bag were put into 3 tubes each containing 1x108 cells. In two tubes, the cells (in volumes of 500-700 µl) were treatment groups, the cells were incubated at 370C for 1 h with 1 µg/ml of each IT, washed and seeded out in soft agar- Assay and incubation (Group A). The cells in the third tube were washed and resuspended to 1 ml in RPMI with 10% FCS (Group B). In the diluted in PBS with 20% ACD and 1% human albumin to a final volume of 1 ml. One of the tubes was used as control and for IT calculations as in Table 1. The plating efficiency in the untreated control cultures was int he range of 20%-30%.

	<del></del>	<del></del>		
rent no. of PBPCs	1 x 108	> >	\ \	
Log tumor cell kill in the presence of different no. of PBPCs	5 x 107	> >	> 5	
Log tumor cell kill i	1 x 107	> 5	> >	
No. of experiments		7	2	
Group Incubation		Unwashed cells treated in normal	saline with ACD Cells washed and	treated in RPMI with 10% FCS
Group		₹	В	

In the apharesis bag the total number of cells will be quite high, and it was conceivable that at such high cell concentrations the efficacy of the procedure might be reduced compared to the conditions used in the model studies. However, in experiments where this possibility was tested no difference in efficacy was observed when the total concentration of cells was increased first from  $1x10^7$  to  $5x10^7$ , and then to  $1x10^8$  per ml (Table 6).

Toxicity of the ITs to Normal Hematopoietic Progenitor Cells. The effect of the ITs on the survival of CFU-GM and BFU-E was studied under incubation conditions as described above. It was found (Table 7) that even a 120 min incubation of the nucleated PBPCs with the IT mixture did not reduce the survival of the progenitor cells, whether tested after washing and resuspending the cells in RPMI with 10% FCS, or when unwashed cells resuspended in normal saline with ACD were used.

Since in a clinical setting the treated cells will be frozen and thawed before given back to the patient, the effect of such procedures on the progenitor cells was also studied. It was found that the freezing and thawing only slightly reduced the number of CFU-GM and BFU-E (Table 7). Notably, the IT treatment itself did not significantly reduce the survival of the progenitor cells, although a slight reduction in the mean number of cell colonies was seen in the group where the cells had been treated under low pH conditions. The data demonstrate that the concentration of ITs that effectively eradicate the tumor cells after 60 min incubation have only insignificant effect on the survival of normal clonogenic cells treated twice as long.

# The state of the s

Table 7

Toxicity of a mixture of MOC-31 and BM7 immunotoxins to CFU-GM and BFU-E in fresh human PBPCs harvested after mobilization with G-CSF. Effect of incubation conditions and the freezing procedure. Control and treatment groups as in Table 6. Nucleated PBPCs were incubated with 1 µg/ml of each of the ITs for 2 h at 370C before being seeded out (5 x 104 cells/dish) in the assay as described in "Materials and Methods".

Incubation		Number of colonies <sup>a</sup> (% of untreated control)	6 of untreated control)	
	CFU	CFU-GM	BFU-E	J-E
	Before freezing	After freezing	Before freezing	After freezing
Untreated	$118 \pm 6 (100)$	102 ± 11 (86)	$170 \pm 15 (100)$	166 ± 56 (98)
Unwashed cells treated in normal saline with 20% ACD	$145 \pm 23 (123)$	86 ± 28 (73)	194 ± 13 (114)	120 ± 42 (71)
Cells washed and treated in RPMI with 10% FCS	125 ± 25 (106)	$112 \pm 16 (95)$	193 ± 73 (114)	159 ± 74 (94)

a Mean ± SD of results obtained from triplicate cultures in two individual experiments

#### DISCUSSION

5

10

15

20

25

30

35

Autologous transplantation of circulating hematopoietic stem cells has recently attracted considerable interest because of its advantages compared to BM transplantation (15, 16). In addition to rapid reconstitution of bone marrow function, it has been supposed that the use of PBSCs might remove the risk of reinfusing tumor cells contaminating the transplant. However, it has been shown that the problem of tumor cell contamination is reduced but not eliminated (4). It should also be noted that high dose chemotherapy involving the use of colony-stimulating factors may recruit tumor cells into the peripheral blood (7, 8). Therefore, a rapid and practical procedure for purging apharesis products is highly warrented.

Several methods for removing breast cancer cells from BM have been reported, including chemo-immunoseparation, immunomagnetic procedures, and the use of immunotoxins (14, 17, 18, 19). In contrast, only very few studies on purging PBSC preparations have been described (20, 21), but ITs prepared with a ribosomeinactivating protein (22, 23) have been used for killing lymphoid tumor cells added to CD34-positive cell collections prepared from BM (24). In the latter study, a purging efficacy of 2 logs was obtained in addition to the 3 log indirect purification achieved by the CD34 selection procedure. The objective of the present study was to develop a safe IT procedure to purge breast cancer cells from PBSCs. The results obtained in model experiments demonstrate that 60 min incubation with 1 μg/ml of each of 2 conjugates involving anti-carcinoma antibodies and PE, efficiently killed all tumor cells admixed to PBSCs without toxicity to the normal progenitor cells. Importantly, the method allows the ITs to be added directly to the apharesis product, and after incubation the cells are washed, centrifuged and ready for freezing. Particularly because of its simplicity and efficacy the method should be attractive for use in the management of selected groups of breast cancer patients in conjunction with high dose chemotherapy combined with transplantation of PBSCs. The high selective efficacy of our procedure might be ascribed to the following factors:

First, the antigen recognized by the MOC31 antibody is known to be expressed on most cells in almost all breast cancer specimens examined (10). Also the BM7 antibody, which recognises the core protein expressed by the MUC-1 gene (9), binds to a high fraction of breast cancer cells (25). Together, these two monoclonals seem to cover, to a reasonable extent, the heterogeneity in antigen expression found in breast cancer. Secondly, we have previously demonstrated that

10

15

20

25

30

when constructing ITs it is important to use a toxin that matches the antibodies used (11). We have found that PE conjugates involving a number of monoclonals, including those that were used here, are very effective (14). Moreover, ITs with PE are always more toxic than equimolar concentrations of free PE (11, 18), demonstrating the specificity of such ITs.

Purging procedures need to be efficient and safe, and it is also necessary that the method is practical and can be used on a clinical scale. In addition to the advantage that the ITs can be added directly to the apharesis bag, our method includes only 60 min incubation time for killing all the clonogenic tumor cells. Moreover, this treatment is not toxic to the normal hematopoietic progenitor cells, and in BM purging experiments even much higher IT concentrations were well tolerated (14). We have also shown that freezing and thawing of the PBSCs treated with the ITs caused no additional toxicity, and it is noteworthy that the IT procedure does not involve the unspecific cell loss that might be experienced with methods involving removal of tumor cells with immunobeads or immunoadsorption.

We have previously calculated the amount of conjugate remaining in IT-treated BM after washing to be about 0.75% of the total amount added (26). In a clinical setting, treatment of PBSCs, containing approximately  $2 \times 10^{10}$  mononuclear cells, with the recommended concentration of 2  $\mu$ g IT/ml ( $1 \times 10^8$  cells), would then be expected to result in a maximum of 3  $\mu$ g IT in the final product. This represents 100-150 times less free toxin than a theoretically calculated maximum tolerable dose (26). Thus, reinfusion of the purged PBSCs would not be expected to give any systemic toxicity.

The success of failure of high dose therapy combined with autologous hematopoietic progenitor cell transplantation may depend even more on the efficacy of the systemic treatment than the efficiency of purging the grafts (1). Nevertheless, it is logical to remove cancer cells that might be present in the autograft, and recent evidence from studies of other tumor types demonstrates the importance of purging (16). In breast cancer, we suggest employing a simple, safe and effective procedure as the one described here.

## Example 2

5

Since breast cancer cells may show different sensitivity to immunotoxins, such that the purging activity of BM7 and MOC31 may be different on breast cancer cells from different patients, the same experiment as previously performed with PM1 breast cancer cells (Example 1) was repeated with another cell line, MA11. It was found that the effect of the immunotoxin treatment was as good, or even better, against the MA11 cells compared to PM1 celles (Table 4). the results confirm the high specific activity of the immunotoxin purging treatment.

In separate experiments the kinetics of the cell killing activity of the immunotoxin was studied in an experiment where PM1 breast cancer cells were added to peripheral blood progenitor cells (ratio 1:100). After incubation for two hours, the mixed cell suspension was frozen and thereafter thawn before the cells were seeded out and the viability of cancer cells and normal progenitor cells assessed in parallell experiments. It was found that intoxication of the breast cancer cells occured rapidly, and within approx. 72 hours all tumor cells were dead. In comparison, no difference in viability of the normal blood progenitor was found in cultures of immunotoxin-treated and non-treated cells within the same time frame.

# 20 Examples 3-4

Carcinoma cells that spread to bone or bone marrow, to pleural and abdominal cavities, to brain and spinal cord tissue, and to the urogenital tract can be selectively killed by immunotoxins administered into the tumor, into said body fluids, or systemically, e.g. to target metastatic tumor cells in tissues such as blood, bone and bone marrow.

#### Example 3

25

30

MA-11 human breast cancer cells were injected into the left cardiac ventricle of immunodeficient rats. Untreated control animals developed symptoms of spinal cord compression and had to be killed 34-37 days after cell injection. Animals treated intravenously with a single dose of MOC31-PE (20 µg/rat) showed a prolonged symptom-free survival and some of the animals lived for more than 50 days.

Another experiment in the same model confirmed the results, and in this case some of the animals survived throughout an observation time of 110 days. In these experiments, one group of rats was treated with an immunotoxin consisting of the 425.3 antibody directed against the EGF-receptor conjugated to PE. All the animals

in this group survived.

In a third experiment in this model, the control rats showed symptoms of cord compression and had to be killed between day 40 and 60 after cell injection. In this experiment three treatment groups were included, one with 20  $\mu$ g 425.3-PE, and one receiving 10 mg cach of the two immunotoxins. Significant prolongation of discase-free survival was obtained with both immunotoxins used individually, giving 60% and 80% long term survival with MOC31-PE and 425.3-PE, respectively. In the combination experiments all animals survived disease-free.

10

15

In a fourth experiment in this model the effect of MOC31-PE was compared to those of cis-platin and doxorubicin. In this experiment all the MOC31-PE treated animals survived for more than 70 days, whereas doxorubicin only showed a marginal effect, and cis-platin-treated rats did not live longer than the saline-treated control animals. These data convincingly demonstrate that the immunotoxins used are highly superior in killing breast cancer metastases than doxorubicin and cisplatin, two of the most commonly used drugs in the clinic.

#### Example 4

- The human breast cancer cell line MT-1 was used in two different experiments. In the first of these, cells were injected into the left cardiac ventricle and the control animals had to be killed because of symptoms of spinal cord compression after a median time of 19 days. Animals treated with 425.3-PE intravenously one day after cell injection all survived. In the other experiments the MT-1 tumor cells were injected directly into the bone marrow cavity of the rat tibia. All untreated animals had to be sacrificed 20 days later because of growing tibial tumors, whereas rats treated with 20 µg of 425.3-PE intravenously one day after cell injection all survived for more than 100 days.
- Furthermore, in the model where MT-1 tumor cells were injected directly into the bone marrow of rat tibia the effect of BM7-PE administered either on day 1 or on day 7 was compared to the effects of 425.3-PE, in groups of animals treated on the same days as for BM7-PE. Morover, the effect of doxorubicin (Adriamycin) given intravenously on day 7 and day 14 was also studied. It was found that both immunotoxins cured 80% of the rats, whether administered on day 1 or on day 7. When half the concentrations of each immunotoxin was combined all animals survived. In comparison, doxorubicin was clearly less effective, leaving only 35% of the animals alive after 90 days. The control animals had to be sacrifized 20 days

after injection of the cells, as in previous experiments. The data confirm the effect of 425.3-PE previously shown. Importantly, BM7-PE was found to be equally effective as 425.3-PE. Both agents were clearly superior in efficacy compared to doxorubicin, one of the clinical drugs most frequently used against breast cancer.

Moreover, the combination of the two immunotoxins cured all animals of their disease.

# Example 5

10

15

20

In two sets of experiments the effect of a recombinantly produced immuntoxin directed against the erbB2-gene product, and with a recombinantly made variant of PE, was tested. In the model described in example 4 the highest concentration of the recombinant immunotoxin significantly prolonged the lifespan of the animals, and 35% of the rats survived. In a model where the MT-1 breast cancer cells were injected intrathecally in immunodeficiant rats, the treatment with the recombinant immunotoxin also given intrathecally (on days 1, 2, and 3) resulted in a significant prolongation in lifespan of the animals. This effect was dose dependent, and the two different doses increased the lifespan of the animals from 10,6 days (saline-treated controls) to 23,4 days and 32,8 days with two different doses of the immuntoxin. At the highest dose 20% of th rats survived. Since no toxicity was observed even with the highest immunotoxin dose, it is expected that the effect at optimal doses may be even better.

#### REFERENCES

5

10

- 1. Peters, W.P., Ross, M., Vredenburgh, J.J., Meisenberg, B., Marks, L.B., Winer, E., Kurtzberg, J., Bast, R.C.J., Jones, R., Shpall, E., Wu, K., Rosner, G., Gilbert, C., Mathias, B., Coniglio, D., Petros, W., Henderson, I.C., Norton, L., Weiss, R.B., Budman, D., and Hurd, D. High-dose chemotherapy and autologous bone marrow support as consolidation after standard-dose adjuvant therapy for high risk primary breast cancer. J. Clin. Oncol., 11: 1132-1143, 1993.
- 2. Armitage, J.O. Bone marrow transplantation. N. Engl. J. Med., 330: 827-838, 1994.
- 3. Moss, T.J., Sanders, D.G., Lasky, L.C., and Bostrom, B. Contamination of peripheral blood stem cell harvests by circulating neuroblastoma cells. Blood, 76: 1879-1883, 1990.
- Ross, A.A., Cooper, B.W., Lazarus, H.M., Mackay, W., Moss, T.J., Ciobanu, N., Tallman, M.S., Kennedy, M.J., Davidson, N.E., Sweet, D., and et al.
   Detection and viability of tumor cells in peripheral blood stem cell collections from breast cancer patients using immunocytochemical and clonogenic assay techniques. Blood, 82: 2605-2610, 1993.
- 5. Brenner, M.K., Rill, D.R., Moen, R.C., Krance, R.A., Mirro, J., Jr., Anderson, W.F., and Ihle, J.N. Gene-marking to trace origin of relapse after autologous bone-marrow transplantation. Lancet, 341: 85-86, 1993.
- Gribben, J.G., Freedman, A.S., Neuberg, D., Roy, D.C., Blake, K.W., Woo, S.D., Grossbard, M.L., Rabinowe, S.N., Coral, F., Freeman, G.J., and et al. Immunologic purging of marrow assessed by PCR before autologous bone marrow transplantation for B-cell lymphoma. N. Engl. J. Med., 325: 1525-1533, 1991.
- 7. Shpall, E.J., and Jones, R.B. Release of tumor cells from bone marrow. Blood, 83: 623-625, 1994.
  - 8. Brugger, W., Bross, K.J., Glatt, M., Weber, F., Mertelsmann, R., and Kanz, L. Mobilization of tumor cells and hematopoietic progenitor cells into peripheral

blood of patients with solid tumors. Blood, 83: 636-640, 1994.

Strous, G.J., and Dekker, J. Mucin-type glycoproteins. Crit. Rev. Biochem. 9. Molecu. Biol., 27: 57-92, 1992.

5

10. Leij, L.D., Postmus, P.E., Poppema, S., Elema, J.D., and The, T.H. The use of monoclonal antibodies for the pathological diagnosis of lung cancer. In: H.H. Hansen (ed), Lung Cancer: Basic and Clinical Aspects, pp. 31-48. Boston: Martinus Niijhoff Publishers, 1986.

10

11. Godal, A., Kumle, B., Pihl, A., Juell, S., and Fodstad, Ø. Immunotoxins directed against the high-molecular-weight melanoma-associated antigen. Identification of potent antibody-toxin combinations. Int. J. Cancer, 52: 631-635, 1992.

15

12. Courtenay, V.D. and Mills, J. An in vitro colony assay for human tumorus grown in immune-suppressed mice and treated in vivo with cytotoxic agents. Br. J. Cancer, 37: 261-268, 1978.

20

35

13. Wang, M.Y., Kvalheim, G., Kvaløy, S., Beiske, K., Jakobsen, E., Wijdens, J., Pihl, A., & Fodstad, Ø. An effective immunomagnetic method for bone marrow purging in T cell malignancies. Bone Marrow Transplant., 9: 319-323, 1992.

14. Myklebust, A.T., Godal, A., Juell, S., Pharo, A., and Fodstad, Ø. Comparison 25 of two antibody-based methods for elimination of breast cancer cells from human bone marrow. Cancer Res., 54: 209-214, 1994.

- 15. Eaves, C.J. Peripheral blood stem cells reach new heights. Blood, 82: 1957-1959, 1993. 30
  - 16. Shpall, E.J., Jones, R.B., Bearman, S.I., Franklin, W.A., Archer, P.G., Curiel, T., Bitter, M., Claman, H.N., Stemmer, S.M., Purdy, M., Myers, S.E., Hami, L., Taffs, S., Heimfeld, S., Hallagan, J., and Berenson, J. Transplantation of enriched CD34-positive autologous marrow into breast cancer patients following high-dose chemotherapy: Influence of CD34-positive peripheralblood progenitors and growth factor on engraftment. J. Clin. Oncol., 12: 28-36, 1994.

15

- 17. Bjorn, M.J., Groetsema, G., and Scalapino, L. Antibody-Pseudomonas exotoxin A conjugates cytotoxic to human breast cancer cells in vitro. Cancer Res., 46: 3262-3267, 1986.
- 18. Anderson, I.C., Shpall, E.J., Leslie, D.S., Nustad, K., Ugelstad, J., Peters, W.P., and Bast, R.C. Jr. Elimination of malignant clonogenic breast cancer cells from human bone marrow. Cancer Res., 49: 4659-4664, 1989.
- 19. O'Briant, K.C., Shpall, E.J., Houston, L.L., Peters, W.P., Bast, R.C. Jr. Elimination of conogenic breast cancer cells from human bone marrow: A comparison of immunotoxin treatment with chemoimmunoseparation using 4-hydroperoxycyclophosphamide, monoclonal antibodies, and magnetic microspheres. Cancer. 68: 1272-1278, 1991.
- Stray, K.M., Corpuz, D., Colter, K.M., Berenson, R., and Heimfeld, S. Purging tumor cells from bone marrow or peripheral blood using avidin biotin immunoadsorptioin. In: P.G. Adrian. G. Samuel, and A.W-W. Diana (eds.), Advances in bone marrow purging and processing, pp. 97-103. Orlando: Wiley-Liss, Inc., 1994.
- Tyer, C.L., Vredenburgh, J.J., Heimer, M., Bast, R.C. Jr., and Peters, W.P.
   Breast cancer cells are effectively purged from peripheral blood progenitor cells using an immunomagnetic technique. Abstract to First meeting of
   International Society for Hematotherapy and Graft Engineering, Orlando, FL, 1993.
- Stirpe, F., Barbieri, L., Battelli, M.G., Soria, M., and Lappi, D.A. Ribosome-inactivating protein from plants: present status and future prospects.
  Bio/Technology 10: 405-412, 1992.
  - 23. Barbieri, L., Battelli, M.G., Stirpe, F. Ribosome-inactivating protein from plants. Biochem. Biophys. Acta. 1154: 237-282, 1993.
- 24. Lemoli, R.M., Tazzari, P.L., Fortuna, A., Bolognesi, A., Gulati, S.C., Stirpe, F., and Tura, S. positive selection of hematopoietic CD34+ stem cells provides 'indirecte purging' of cd34-lymphoid cells and the purging effeciency is increased by anti-CD2 and CD30 immunotoxins. Bone Marrow Transplant.,

10

13: 465-471, 1994.

- 25. Diel, I.J., Kaufmann, M., Goerner, R., Costa, S.D., Kaul, S., and Bastert, G. Detection of tumor cells in bone marrow of patients with primary breast cancer: a prognostic factor for distant metastasis. J. Clin. Oncol., 10: 1534-1539, 1992.
- 26. Myklebust, A.T., Godal, A., Pharo, A., Juell, S., and Fodstad, Ø. Eradication of small cell lung cancer cells from human bone marrow with immunotoxins. Cancer Res., 53: 3784-3788, 1993.



#### CLAIMS

Method to kill breast cancer cells or other carcinoma cells expressing the same target antigens in a cell population comprising nucleated cells harvested from peripheral blood, or CD-34+ cells selected from the above nucleated cells, or other immature/early progenitor cells from blood containing multipotent stem cells, characterized in that the cell population is exposed to a combination of two immunotoxins, wherein each immunotoxin is composed of a conjugate between an antibody and a cell toxin, fragments of antibodies and toxin, or recombinantly produced antibodies, toxins, immunotoxins or fragments thereof, wherein the antibodies are directed to epitopes on the antigen EGP2 expressed by the gene GA733 and to epitopes on the antigen expressed by the genes MUC1, MUC2 or MUC3, respectively or a combination of these, and the toxin is Pseuctomonas exotoxin A.

15

2. Method according to claim 1, characterized in that the used antibodies are MOC31 and an antibody directed to the antigens encoded by the genes MUC1, MUC2, MUC3 or a combination of

these.

20

- 3. Method according to claims 1, 2, characterized in that the used antibodies are MOC31 and BM7, or fragments thereof.
- 4. Method according to claims 1, 2, characterized in that the used anithodies are MOC31 and BM2 or 12H12, or fragments thereof.
  - 5. Method according to claims 1, 2,
- characterzied in that the used antibodies are MOC31 and 595A6, or fragments thereof
  - 6. Method according to claim 1, characterized in that the specific immunotoxins are administered in vivo.
- 35
- 7. Method according to claim 6, characterized in that the immunotoxins are administered systemically, especially in case of malignant spread to tissues such as bone and bone marrow.

- 8. Method according to claim 6, characterized in that the immunotoxins are administered directly into the tumor or in the pleural and abdominal cavities.
- 5 9. Immunotoxins to kill cells according to the method in claim 1, characterized in that it contains two immunotoxins directed to antigens present on malignant cells.
  - 10. Immunotoxins according to claim 9,
- characterized in that the antibodies are selected among MOC31, and BM7, 595, BM2, 12H12 or combinations of these or fragments thereof and the toxin is native or recombinant Pseudomonas exotoxin A, or fragments thereof.
- 11. Use of the immunotoxins according to claim 9, to produce a therapeutic agent against cancer.
  - 13. Kit to perform the method according to claim 1, characterized in that it contains a preparation of two immunotoxins in a pharmaceutically acceptable formulation.

ttorney Docket No. 7885.55USWO

## MERCHANT, GOULD, SMITH, EDELL, WELTER & SCHMIDT

#### **United States Patent Application**

#### COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor I hereby declare that: my residence, post office address and citizenship are as stated below next to my ame; that

I verily believe I am the original, first and sole inventor (if only one name is listed below) or a joint inventor (if pharal inventors are amed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: METHOD OF KILLING ARGET CELLS IN HARVESTED CELL POPULATIONS WITH ONE OR MORE IMMUNOTOXINS he specification of which is attached hereto (if applicable) (in the case of a PCT-filed application) described and was amended on . 🛛 was filed on as application serial no. nd claimed in international no. PCT/NO97/00074 filed Intl Filing Date March 12, 1997 and as amended on (if any), which I have eviewed and for which I solicit a United States patent. hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by ny amendment referred to above. acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of ederal Regulations, § 1.56 (attached hereto). hereby claim foreign priority benefits under Title 35, United States Code, § 119/365 of any foreign application(s) for patent or inventor's crtificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before hat of the application on the basis of which priority is claimed: . no such applications have been filed. . X such applications have been filed as follows: FOREIGN APPLICATION(S), IF ANY, CLAIMING PRIORITY UNDER 35 USC | 119 DATE OF ISSUE DATE OF FILING APPLICATION NUMBER COUNTRY (day, month, year) (day, month, year) 961031 March 13, 1996 Norway ALL FOREIGN APPLICATION(S), IF ANY, FILED BEFORE THE PRIORITY APPLICATION(S) DATE OF FILING DATE OF ISSUE APPLICATION NUMBER COUNTRY (day, month, year) (day, mosth, year) I hereby claim the benefit under Title 35, United States Code, § 120/365 of any United States and PCT international application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37. Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application. STATUS (patented, pending, abandoned) DATE OF FILING (day, month, year) U.S. APPLICATION NUMBER I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below: DATE OF FILING (Day, Month, Year) U.S. PROVISIONAL APPLICATION NUMBER

#### INDEPENDENT INVENTOR(S)

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS (37 C.F.R. 1.9(f) AND 1.27(b)) - INDEPENDENT INVENTOR

As a below named inventor, I hereby declare that I qualify as an independent inventor as defined in 37 C.F.R. 1.9(c) for purposes of paying reduced fees under Section 41(a) and (b) of Title 35, United States Code, to the Patent and Trademark Office with regard to the invention entitled METHOD OF KILLING TARGET CELLS IN HARVESTED CELL POPULATIONS WITH ONE OR MORE IMMUNOTOXINS described in

b) □ I	he specification filed he provisional application s non-provisional application application in the provisional application is the provisional application in the provision in the provisional application in the provision in the provisi	erial no, filed tion serial no. <u>PCT/NO97/00074</u> , filed <u>Ir</u>	atl Filing Date March 12, 1997.
convey or license under 37 C.F.R.	e, any rights in the invent 1.9(c) if that person has	r licensed and am under no obligation un- tion to any person who could not be class made the invention, or to any concern wi or a nonprofit organization under 37 C.F	sified as an independent inventor nich would not qualify as a small
Each person, con under contract or	cern or organization to law to assign, grant, co	which I have assigned, granted, conveyed nvey, or license any rights in the invention	l, or licensed or am under obligation on is listed below:
	a) no such person, b) persons, concer	, concern, or organization ns or organizations listed below*	
	named person	arate verified statements are required fro	o the
ADDRESS 1	Oystein Fodstad Frits Kiacrs v. 28, 0383 a) MINDIVIDUAL	Oslo Norway b) SMALL BUSINESS CONCERN c)	□ NONPROFIT ORGANIZATION
NAME ADDRESS	a)   INDIVIDUAL	b) SMALL BUSINESS CONCERN c)	□ NONPROFIT ORGANIZATION
entitlement to sn	nall entity status prior to	plication or patent, notification of any ch paying, or at the time of paying, the earl sich status as a small entity is no longer a	iest of the issue fee or any
information and willful false state	belief are believed to be ements and the like so m inited States Code, and i	e herein of my own knowledge are true as true; and further that these statements we hade are punishable by fine or imprisonm that such willful false statements may jeour any patent to which this verified statem	ere made with the knowledge that ent, or both under Section 1001 of pardize the validity of the
Oystein Fodstad		Gunnar Kvalheim	Meng yu Wang NAME OF INVENTOR
NAME OF INV	ENTOR Les Took	MAME OF INVENTOR	Margyll Wary
Signature of Inv	rentor 18 - 28	Signature of Inventor	Signature of Inventor
Date	17-00	Date	Date

#### INDEPENDENT INVENTOR(S)

# VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS (37 C.F.R. 1.9(f) AND 1.27(b)) - INDEPENDENT INVENTOR

As a below named inventor, I hereby declare that I qualify as an independent inventor as defined in 37 C.F.R. 1.9(c) for purposes of paying reduced fees under Section 41(a) and (b) of Title 35, United States Code, to the Patent and Trademark Office with regard to the invention entitled <u>METHOD OF KILLING TARGET CELLS IN HARVESTED CELL POPULATIONS WITH ONE OR MORE IMMUNOTOXINS</u> described in

	b) [		n serial no, filed ication serial no. <u>PCT/NO97/0</u>	0074, filed Intl Filing Date Marc	h 12, 1997.
	convey or licer under 37 C.F.I	nse, any rights in the inv R. 1.9(c) if that person h	ention to any person who cou	obligation under contract or law to describe the classified as an independence of the concern which would not qualify under 37 C.F.R. 1.9(e).	dent inventor
30, 37				ted, conveyed, or licensed or am named in the invention is listed below:	under obligation
Species The World		a) no such pers b) persons, con	on, concern, or organization cerns or organizations listed b	elow*	
Ber and Roun and		named pers	eparate verified statements are son, concern or organization he verring to their status as small	aving rights to the	
The Section	NAME ADDRESS	Oystein Fodstad Frits Kiaers v. 28, 038 a) ⊠ INDIVIDUAL	3 Oslo Norway  b) SMALL BUSINESS C	ONCERN c)   NONPROFIT OR	GANIZATION
The first for the	NAME ADDRESS	a)   INDIVIDUAL	b) ☐ SMALL BUSINESS C	ONCERN c)   NONPROFIT OR	GANIZATION
	I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 C.F.R. 1.28(b))				
	I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereof, or any patent to which this verified statement is directed.				
	Olav Engebrai		Siri Juell NAME OF INVENTOR	NAME OF BUVE	AFTOD
	NAME OF IN	VENTOR	Sui Tuell	NAME OF INVE	
•	Signature of I	rventor	Signature of Inventor	Signature of Inver	itor
•	Date 24	4/28	Date	Date	

I hereby appoint the following attorney(s) and/or patent agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected herewith:

mark critica confection notewith;			
Albrecht, John W.	Reg. No. 40,481	Lacy, Paul E.	Reg. No. 38,946
Ansems, Gregory M.	Reg. No. P-42,264	Larson, James A.	Reg. No. 40,443
Batzli, Brian H.	Reg. No. 32,960	Lasky, Michael B.	Reg. No. 29,555
Beard, John L.	Reg. No. 27,612	Lindquist, Timothy A.	Reg. No. 40,701
Berman, Charles	Reg. No. 29,249	Lynch, David W.	Reg. No. 36,204
Black, Bruce E.	Reg. No. 41,622	Marschang, Diane L.	Reg. No. 35,600
Blasdell, Thomas L.	Reg. No. 31,329	McDaniel, Karen D.	Reg. No. 37,674
Bogucki, Raymond A.	Reg. No. 17,426	McDonald, Daniel W.	Reg. No. 32,044
Bruess, Steven C.	Reg. No. 34,130	McIntyre, Iain A.	Reg. No. 40,337
Byrne, Linda M.	Reg. No. 32,404	Mueller, Douglas P.	Reg. No. 30,300
Canady, Karen S.	Reg. No. 39,927	Nasiedlak, Tyler L.	Reg. No. 40,099
Carlson, Alan G.	Reg. No. 25,959	Nelson, Albin J.	Reg. No. 28,650
Carter, Charles G.	Reg. No. 35,093	Pauly, Daniel M.	Reg. No. 40,123
Caspers, Philip P.	Reg. No. 33,227	Plunkett, Theodore	Reg. No. 37,209
Chiapetta, James R.	Reg. No. 39,634	Pytel, Melissa J.	Reg. No. 41,512
Clifford, John A.	Reg. No. 30,247	Reich, John C.	Reg. No. 37,703
Daignault, Ronald A.	Reg. No. 25,968	Reiland, Earl D.	Reg. No. 25,767
Daley, Dennis R.	Reg. No. 34,994	Rittmaster, Ted R.	Reg. No. 32,933
Dalglish, Leslie E.	Reg. No. 40,579	Schmaltz, David G.	Reg. No. 39,828
Daulton, Julie R.	Reg. No. 36,414	Schuman, Mark D.	Reg. No. 31,197
DeVries Smith, Kate	Reg. No. P-42,157	Schumann, Michael D.	Reg. No. 30,422
DiPietro, Mark J.	Reg. No. 28,707	Sebald, Gregory A.	Reg. No. 33,280
Edell, Robert T.	Reg. No. 20,187	Skoog, Mark T.	Reg. No. 40,178
Epp Ryan, Sandra	Reg. No. 39,667	Soderberg, Richard	Reg. NoP-43,352
Farber, Michael B.	Reg. No. 32,612	Sumner, John P.	Reg. No. 29,114
Funk, Steven R.	Reg. No. 37,830	Sumners, John S.	Reg. No. 24,216
Glance, Robert J.	Reg. No. 40,620	Tellekson, David K.	Reg. No. 32,314
Golla, Charles E.	Reg. No. 26,896	Trembath, Jon R.	Reg. No. 38,344
Gorman, Alan G.	Reg. No. 38,472	Underhill, Albert L.	Reg. No. 27,403
Gould, John D.	Reg. No. 18,223	Vandenburgh, J. Derek	Reg. No. 32,179
Gregson, Richard	Reg. No. P-41,804	Vradenburgh, Anna M.	Reg. No. 39,868
Gresens, John J.	Reg. No. 33,112	Welter, Paul A.	Reg. No. 20,890
Hamre, Curtis B.	Reg. No. 29,165	Whipps, Brian	Reg. No. P-43,261
Hillson, Randall A.	Reg. No. 31,838	Williams, Douglas J.	Reg. No. 27,054
Johnston, Scott W.	Reg. No. 39,721	Witt, Jonelle	Reg. No. 41,980
Kastelic, Joseph M.	Reg. No. 37,160	Wood, Gregory B.	Reg. No. 28,133
Kettelberger, Denise	Reg. No. 33,924	Wood, William J.	Reg. No. P-42,236
Komanduri, Janaki	Reg. No. 40,684	Xu, Min S.	Reg. No. 39,536
Kowalchyk, Alan W.	Reg. No. 31,535		African cognitation African communication co
Kowalchyk, Katherine M.	Reg. No. 36,848		

I hereby authorize them to act and rely on instructions from and communicate directly with the person/assignee/attorney/firm/ organization who/which first sends/sent this case to them and by whom/which I hereby declare that I have consented after full disclosure to be represented unless/until I instruct Merchant, Gould, Smith, Edell, Welter & Schmidt to the contrary.

Please direct all correspondence in this case to Merchant, Gould, Smith, Edell, Welter & Schmidt at the address indicated below:

Merchant, Gould, Smith, Edell,

Welter & Schmidt

3100 Norwest Center

90 South Seventh Street

Minneapolis, MN 55402-4131

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

2	Full Name Of Inventor	Family Name  Fodstad  OO	First Given Name / Oystein		Second Given Name
0	Residence & Clitzenship	City Onlo	State or Foreign Country Norway		Country of Citizenship Norway
1	Post Office Address	Post Office Address Fritz Kinors v. 28	City Oslo -		State & Zip Code/Country Norway 0323
Signs	ture of Inventor 2			Date:	23/9-98
2	Full Name Of Inventor	Family Name  Kvalheim  2 00	First Given Name Ounnar		Second Given Name
•	Residence & Citizenship	City NO	State or Foreign Country Norway		Country of Citizenship Norway
2	Post Office Address	Post Office Address Asstubben 13	City Oslo		State & Zip Code/Country N=0381 Norway
Signa	ture of Inventor 2	Dunnal Walkin		Date:	25/4-99
2	Full Name Of Inventor	Family Name (Wang 3-60)	First Given Name Meng		Second Given Name yu
•	Residence & Citizenship	City NO X	State or Foreign Country Norway		Country of Citizenship China
3	Post Office Address	Post Office Address Montsbellovn 11	City Oslo		State & Zip Code/Country Norway N=0379
Signature of Inventor 203:				Date:	29.09.98
2	Full Name Of Inventor	Family Name /Engebraten / 00	First Given Name Olav		Second Given Name
•	Residence & Citizenship	City Lorgiskog	State or Foreign Country Norway		Country of Cittzenskip Norway
4	Post Office Address	Post Office Address Haneborgsvn 43 A	City Lorenskog		State & Zip Code/Country N-1470 Norway
Signature of Inventor 204: Olen Tylica. Date:					28/5-98
2	Full Name Of Inventor	Family Name Juell 5-00	First Gives Name		Second Given Name
0	Residence & Citizenship	City NOX	State or Foreign Country Norway		Country of Citizenship Norway
5	Post Office Address	Post Office Address Granholtet 6	City Lommedalen		State & Zip Code/Country N-1350 Norway
Sign	ature of Inventor 2	Date:	7.10.98		

#### § 1.56 Duty to disclose information material to patentability.

- (a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is canceled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is canceled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability is deemed to be satisfied if all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)–(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced or attempted or the duty of disclosure was violated through bad faith or intentional misconduct. The Office encourages applicants to carefully examine:
  - (1) prior art cited in search reports of a foreign patent office in a counterpart application, and
- (2) the closest information over which individuals associated with the filing or prosecution of a patent application believe any pending claim patentably defines, to make sure that any material information contained therein is disclosed to the Office.
- (b) Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and
  - (1) It establishes, by itself or in combination with other information, a prima facie case of unpatentability of a claim;

or o

- (2) It refutes, or is inconsistent with, a position the applicant takes in:
  - (i) Opposing an argument of unpatentability relied on by the Office, or
  - (ii) Asserting an argument of patentability.

A prima facie case of unpatentability is established when the information compels a conclusion that a claim is unpatentable under the preponderance of evidence, burden—of—proof standard, giving each term in the claim its broadest reasonable construction consistent with the specification, and before any consideration is given to evidence which may be submitted in an attempt to establish a contrary conclusion of patentability.

- (c) Individuals associated with the filing or prosecution of a patent application within the meaning of this section are:
  - (1) Each inventor named in the application:
  - (2) Each attorney or agent who prepares or prosecutes the application; and
- (3) Every other person who is substantively involved in the preparation or prosecution of the application and who is associated with the inventor, with the assignee or with anyone to whom there is an obligation to assign the application.
- (d) Individuals other than the attorney, agent or inventor may comply with this section by disclosing information to the attorney, agent, or inventor.